

The Microbiological Hazards of *Enterobacter* spp. Contamination in Street Food in Developing Countries: A Literature Review

Jefyne Mali' Pareakan¹, Charis Amarantini¹, Tri Yahya Budiarto^{1*}

¹ Biology Study Program, Faculty of Biotechnology, Universitas Kristen Duta Wacana, Yogyakarta, Indonesia

*Corresponding Author Email: yahya@staff.ukdw.ac.id

Copyright: ©2025 The author(s). This article is published by Media Publikasi Cendekia Indonesia.

LITERATURE REVIEW

Submitted: 10 May 2025

Accepted: 2 June 2025

Keywords:

Developing countries, *Enterobacter*, Foodborne, Food safety, Microbiological hazard, Street food

OPEN ACCESS



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License

ABSTRACT

Street food is widely consumed by people in developing countries and often poses health risks due to bacterial contamination. The consumption of contaminated food has been linked to more than 200 types of diseases, resulting in 600 million cases and 420,000 deaths each year. One commonly found bacterium is *Enterobacter*, which can be pathogenic and exhibit resistance to various antibiotics. This review aims to assess the dangers of *Enterobacter* contamination in street food, including the level and source of contamination, pathogenicity, and management strategies. The review followed PRISMA 2020 guidelines, with literature searches performed in scientific databases such as PubMed and Google Scholar using Boolean operators and relevant keywords. Articles were selected based on specific inclusion criteria, analyzed, and grouped for ease of interpretation. The results showed that *Enterobacter* contamination is prevalent in developing countries such as Nigeria, Ghana, Ethiopia, Pakistan, India, and Indonesia. Nigeria recorded the highest prevalence, with *E. aerogenes* reaching 100% in ready-to-eat rice. The primary source of contamination is raw materials, particularly wheat flour, which contains *E. sakazakii* with a prevalence of 100%. *Enterobacter* can cause various serious infections, including urinary tract infections, meningitis, pneumonia, and septicemia. Efforts to reduce *Enterobacter* contamination in street food should include the implementation of hygiene and sanitation practices, the use of safe raw materials, proper processing and storage methods, education for vendors, and routine food safety supervision. These findings are crucial for supporting public health interventions and developing effective food safety policies.

Key Messages:


- The presence of *Enterobacter* in street food represents a potential public health concern, as it may contribute to foodborne illnesses and nosocomial infections due to its marked pathogenicity, virulence factors, and substantial antibiotic resistance.
- Contamination of street food with *Enterobacter* is commonly caused by poor hygiene and sanitation practices, underscoring the need to implement effective food safety management systems to prevent the risk of contamination.

Access this article online



Quick Response Code

GRAPHICAL ABSTRACT



INTRODUCTION

The global consumption of food contaminated with pathogenic bacteria is increasing, leading to more than 200 types of potentially fatal infections, particularly affecting infants, the elderly, and pregnant women (1,2). This issue impacts approximately 600 million people about 1 in 10 of the global population and results in an estimated 420,000 deaths annually (3). Foodborne infections are primarily associated with the consumption of street food, ready-to-eat (RTE) food, and fast food (4,5). Street food refers to food sold outdoors, in public places, or on the roadside, prepared and served directly for consumption without the need for further processing (6,7). The development of street food sector, which operates without adequate regulations regarding hygiene and safety standards increases the risk of microbiological contamination. Street food vendors face several challenges in maintaining food hygiene and safety, including limited access to clean water, the use of low-quality raw materials, improper packaging methods, inappropriate storage temperatures, and unsupportive environmental conditions (6). These factors contribute to rapid bacterial growth, thereby elevating the risk of foodborne infections. For instance, a listeriosis outbreak in South Africa in 2018 infected 674 people and caused 183 deaths (8). Similarly, in Bangladesh, approximately 30 million people contract foodborne infections annually, with an estimated 2.2 million dying from diarrhea (5).

Escherichia coli and *Enterobacter* are the most common pathogenic bacteria found in street food (9,10). A study conducted in Nigeria revealed that 50% of street food samples were contaminated with *Enterobacter* spp. (11), while in India, 36.7% of 44 samples were infected (12), and in Ethiopia, 15.87% of 44 samples also showed similar contamination (13). *Enterobacter* can produce various cellular toxins, including enterotoxins, hemolysins, and pore-forming toxins, which significantly contribute to foodborne diseases (14). Studies indicate that the types of *Enterobacter* commonly found in street food include *E. cloacae*, *E. aerogenes*, *E. agglomerans*, and *E. Hafniae* (15,16). Although numerous studies have examined microbial contamination in street food, there has been limited focus on the hazards posed by *Enterobacter* contamination in developing countries. Previous research has emphasized pathogens such as *Escherichia coli*, *Salmonella*, and *Staphylococcus aureus*. It has been demonstrated that *Enterobacter* contamination can lead to nosocomial infections and gastrointestinal diseases through the fecal-oral route, presenting symptoms such as fever, diarrhea, and abdominal pain (15,17). Therefore, there is a pressing need for more in-depth studies on the microbiological hazards associated with *Enterobacter* contamination in street

food within developing countries. This literature review aims to highlight the dangers related to *Enterobacter* contamination in street food, focusing on prevalence, sources, risk factors, and its impact on public health. It is expected that this review will significantly contribute to the development of evidence-based food safety policies, the improvement of hygiene practices in food handling, and the formulation of public health risk mitigation strategies concerning *Enterobacter* spp. contamination in street food in developing countries.

METHODS

This review was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses 2020 (PRISMA) guidelines.

Search strategy

Data collections were performed by searching scientific articles across multiple databases, including Google Scholar and ScienceDirect. The search employed Boolean operators ("AND," "OR," "NOT") combined with keywords relevant to the study topic, such as *Enterobacter*, street food or ready-to-eat food, microbial contamination, foodborne infection, and developing countries. Various keyword combinations were used to maximize the retrieval of pertinent articles. The search and article selection process were systematically documented using a PRISMA flow diagram (Figure 1), illustrating the number of articles identified, screened, assessed for eligibility, and included in the analysis.

Inclusion and Exclusion Criteria

This review included studies conducted in various countries and comprised original research articles, review articles, peer-reviewed papers, and systematic reviews. Inclusion criteria were: articles published in English, available in full text, and published between 2010 and 2025. Selected articles had to be relevant to the topic, particularly those reporting data on the prevalence of contamination and microbiological hazards posed by bacteria, especially *Enterobacter* species, in street food within developing countries. Articles providing information on the pathogenic characteristics, virulence mechanisms, and types of infections or diseases caused by *Enterobacter* at the individual and population levels were also included. Exclusion criteria encompassed editorials, opinion pieces, articles unavailable in full text, and studies not focused on human health.

Study selection

The article selection process was conducted in multiple stages. Initially, all retrieved articles were de-duplicated using reference management software (Zotero). Subsequently, titles and abstracts were screened to assess their preliminary eligibility based on the inclusion criteria.

Data extraction and analysis

Data were systematically extracted from articles that met the inclusion criteria using a pre-designed data extraction form. The form covered three main aspects: (1) levels of *Enterobacter* contamination in various developing countries, (2) sources of *Enterobacter* contamination in street food, and (3) types of infections caused by *Enterobacter* species. Extracted information from each article included the author's name, year of publication, study location, type of food tested, identified *Enterobacter* species, prevalence rates, contamination sources, and reported health effects or infections.

Descriptive and thematic analyses were conducted. Data were categorized and presented in three separate tables to facilitate visualization and interpretation. Thematic analysis was employed to identify common patterns, regional differences, and potential public health risks associated with the presence of *Enterobacter* in street food. The synthesized findings formed the basis for drawing conclusions and making recommendations aimed at controlling microbiological risks in street food within developing countries.




Figure 1. Prisma Flow Diagram

RESULTS

Contamination Profiles of Pathogenic Bacteria in Street Food across Different Countries

The levels of pathogenic bacterial contamination in street food vary significantly across countries, reflecting differences in sanitation practices, food processing, and handling methods. Table 1 summarizes the findings by geographical region and type of food sample analyzed, providing a comprehensive overview of the prevalence of pathogenic bacterial contamination in street foods worldwide.

Table 1. Contamination levels of *Enterobacter* and other pathogenic bacteria in street food in developing countries

Country	Types of food	Prevalence of <i>Enterobacter</i>	Prevalence of other bacteria	References
Nigeria	RTE-cooked rice	<i>E. aerogenes</i> (100%)	<i>S. aureus</i> (91.7%) <i>Brahmella</i> sp. (80.6%) <i>Salmonella</i> spp. (69%)	(18)
Nigeria	Rice, soup, and beans	<i>Enterobacter</i> spp.	<i>E. coli</i> (%)	(11)

Country	Types of food	Prevalence of <i>Enterobacter</i> (50%)	Prevalence of other bacteria	References
			<i>Streptococcus spp.</i> (50%) <i>S. aureus</i> (60%) <i>P. aeruginosa</i> (90%) <i>Salmonella spp.</i> (30%) <i>B. cereus</i> (40%) <i>Micrococcus spp.</i> (30%) <i>A. faecalis</i> (10%)	
Africa	Vegetables, potatoes, rice, pies, beef, and chicken stew	<i>Enterobacter spp.</i> (18%)	<i>Listeria spp.</i> (22%) <i>A. hydrophila</i> (12%) <i>K. oxytoca</i> (8%) <i>P. mirabilis</i> (6.3%) <i>S. aureus</i> (3.2%) <i>P. luteola</i> (2.4%)	(19)
Ghana	Ampesi, banku, beans, bofloat, bread, fried egg, fried fish/ meat, fried, pepper, fried rice, gari, ground pepper, hausa kooko, indomie/ spaghetti, jollof, kenkey, rice, salad, soup, stew, waakye	<i>Enterobacter spp.</i> (16.8%)	<i>Citrobacter spp.</i> (10.1%) <i>E. faecalis</i> (7.8%) <i>Pseudomonas spp.</i> (6.7%) <i>K. pneumoniae</i> (4.0%)	(10)
Ethiopia	Sanbusa, Donat, Bombolino, and bread	<i>Enterobacter spp.</i> (15.87%)	<i>S. aureus</i> (53.96%) <i>E.coli</i> (23.8%) <i>Citrobacter spp.</i> (6.3%)	(13)
Pakistan	Nehari, wonton, wonton, chicken shashlik, grilled chicken, stuffed chicken, pani puri, haleem, chicken manchurian, chicken mayo garlic roll, sandwich, french fries, bbq items, zinger, samosa, coleslaw salad, finger fish	<i>E. aerogenes</i> (10%)	<i>Pseudomonas spp.</i> (20%) <i>Bacillus spp.</i> (18%), <i>S. aureus</i> (6%) <i>S. epidermidis</i> (6%) <i>S. fecalis</i> (2%) <i>E. coli</i> (14%)	
India	Panipuri, chaats, and eggrolls	<i>Enterobacter spp.</i> (8,93%)	<i>E. coli</i> (37.5%) <i>P. aeruginosa</i> (3.57%) <i>S. aureus</i> (14.20%) <i>Salmonella spp.</i> (5.36%) <i>Klebsiella spp.</i> (10.71%) <i>Shigella spp.</i> (19.64%)	(20)
Indonesia	Cilok, dumplings, and skewer meatballs	<i>E. cloacae</i> (6.6%)	<i>E. coli</i> (16.6%) <i>K. pneumoniae</i> (13.3%) <i>Y. enterocolitica</i> (13.3%) <i>Pantoea spp.</i> (6.6%) <i>A. hydrophila</i> (6.6%) <i>S. marcescens</i> (6.6%) <i>Moraxella spp.</i> (6.6%) <i>S. liquefaciens</i> (3.3%) <i>P. mirabilis</i> (3.3%) <i>Shigella spp.</i> (3.3%) <i>E. Americana</i> (3.3%)	(21)
India	Fried rice, jollof rice, moi-moi, salad, oil beans, non-oil beans, and African salad	<i>E. cloacae</i> (5.36%)	<i>E. coli</i> (41.07%) <i>C. freundii</i> (5.36%)	(16)

The prevalence of *Enterobacter* genus bacteria in street food from developing countries showed considerable variation, influenced by the type of food as well as local environmental and sanitation conditions. Notably, the highest prevalence of *Enterobacter aerogenes* was observed in ready-to-eat (RTE) cooked rice in Nigeria, reaching 100%. In contrast, countries such as Ghana, Ethiopia, Pakistan, India, and Indonesia reported *Enterobacter* spp. prevalence rates ranging from 5.36% to 50%. In addition to *Enterobacter*, other predominant pathogenic bacterial isolates identified included *Staphylococcus aureus*, *Escherichia coli*, *Salmonella* spp., and *Pseudomonas* spp. These findings suggest potential cross-contamination and multispectral infection risks for consumers. Overall, these results underscore the urgent need to strengthen microbiological surveillance systems and enhance hygienic practices in street food preparation and handling to minimize pathogen exposure and mitigate associated public health risks.

Source of *Enterobacter* Contamination in Street Food

The presence of bacterial contamination in street food is influenced by various factors, including the quality of raw materials, environmental cleanliness, hygienic practices of food handlers, and food processing and storage methods. These factors interact synergistically, increasing the risk of bacterial contamination and proliferation on food, which may lead to adverse health effects for consumers. Table 2 presents the identified sources of bacterial contamination in street food.

Table 2. Sources of *Enterobacter* contamination and other pathogenic bacteria in street food

Source	Prevalence of <i>Enterobacter</i>	Prevalence of Other Bacteria	References
Wheat flour	<i>E. sakazakii</i> (100%)	-	(22)
Raw vegetable	<i>Enterobacter</i> spp. (64%)	<i>S. aureus</i> (48%) <i>P. aeruginosa</i> (60%) <i>E. coli</i> (48%) <i>Salmonella</i> spp. 8%	(23)
Fresh Vegetables	<i>E. cloacae</i> (61,46%) <i>E. aerogenes</i> (2,08%) <i>E. cancerogenus</i> (2,08%) <i>E. gergoviae</i> (1,04%) <i>E. Sakazakii</i> (4,17%)	<i>K. oxytoca</i> (19,79%) <i>K. pneumoniae</i> (9,38%)	(24)
RTE/slice fruit	<i>Enterobacter</i> (40%)	<i>Bacillus</i> spp. (100%) <i>S. aureus</i> (80%) <i>Penicillium</i> spp. (80%) <i>A. niger</i> (60%) <i>E. coli</i> (40%) <i>Salmonella</i> (40%) <i>Klebsiella</i> (40%) <i>Mucor</i> spp. (40%) <i>P. aeruginosa</i> (20%) <i>Proteus</i> (20%) <i>Micrococcus</i> (20%) <i>Lactobacillus</i> spp. (20%)	(25)
Dishwashing water	<i>Enterobacter</i> spp. (36.67%)	<i>E. coli</i> (50%) <i>P. aeruginosa</i> (26.67%)	(12)
RTE meat beef, pork, and chicken	<i>E. cloacae</i> (29%)	<i>Escherichia</i> (27%) <i>P. mirabilis</i> (25%)	(26)
RTE meat chicken and beef	<i>E. cloacae</i> (27%) <i>E. sakazakii</i> (13%) <i>E. kobei</i> (5%) <i>E. ludwigii</i> (8%) <i>E. hormaechei</i> (4%)	<i>P. luteola</i> (5%) <i>P. aeruginosa</i> (4%) <i>P. stutzeri</i> (4%) <i>K. pneumonia</i> (15%) <i>K. oxytoca</i> (14%) <i>C. koseri</i> (2%)	(27)
Raw fruits and vegetables	<i>Enterobacter</i> spp. (25,3%)	<i>E. coli</i> (17,4%) <i>Klebsiella</i> spp. (43%) <i>Proteus</i> spp. (4%) <i>Citrobacter</i> spp. (2%)	(28)

Source	Prevalence of <i>Enterobacter</i>	Prevalence of Other Bacteria	References
		<i>Providencia spp.</i> (0,7%)	
Cooking water, storage water, and washing water	<i>E. cloacae</i> (11.1 %).	<i>K. pneumoniae</i> (20.8 %) <i>Aeromonas spp.</i> (16.7 %)	(10)
Kitchen instruments	<i>Enterobacter</i> (9,40%)	<i>S. aureus</i> (15.05%) <i>S. epidermidis</i> (16.50%) <i>E. coli</i> (18.27%) <i>Shigella spp.</i> (0.91%) <i>Bacillus spp.</i> (13.02%) <i>Pseudomonas</i> (2.11%)	(29)
Hand rinse	<i>Enterobacter spp.</i> (9.1%)	<i>S. aureus</i> (23.5%) <i>Klebsiella spp.</i> (16.1%) <i>E. coli</i> (10.9%) <i>Citrobacter spp.</i> (4.3%) <i>S. marcescens</i> (2.6%) <i>P. aeruginosa</i> (3.5%) <i>Proteus spp.</i> (2.2%) <i>P. rettgeri</i> (1.3%) <i>Salmonella spp.</i> (0.9%)	(30)
Pupuru and plantain flour	<i>Enterobacter spp.</i> (0.99%)	<i>Klebsiella spp.</i> (91.12%) <i>Acinetobacter sp.</i> (1.97%) <i>Campylobacter sp.</i> (4.61%) <i>Corynebacterium sp.</i> (0.33%) <i>B. subtilis</i> (0,99%)	(31)

Data in Table 2 indicate that *Enterobacter* species are widespread across various food items and environments associated with street food. High prevalence rates were observed in wheat flour (*E. sakazakii*, 100%), fresh vegetables (*E. cloacae*, 61.46%), and ready-to-eat cut fruits (40%). Significant contamination was also detected in ready-to-eat processed meats such as chicken and beef (*E. cloacae*, 27–29%), as well as in environmental sources, including equipment wash water (36.67%) and kitchen utensils (9.4%). These contaminations frequently co-occur with other pathogenic bacteria such as *Escherichia coli*, *Staphylococcus aureus*, *Klebsiella spp.*, and *Salmonella spp.*, cumulatively heightening the risk of transmission of gastrointestinal infections and other opportunistic diseases.

Infections Caused by *Enterobacter*

Contamination of street food by *Enterobacter* species poses a potential risk of infection to consumers, as these bacteria are opportunistic pathogens capable of causing a variety of nosocomial and community-acquired infections. Table 3 summarizes the types of infections caused by different *Enterobacter* species, highlighting their clinical significance and public health impact.

Table 3. Various types of Infections caused by various *Enterobacter* species

<i>Enterobacter</i>	Infection	Source
<i>E. cloacae</i>	Endocarditis, septic arthritis, osteomyelitis, urinary tract infections (UTIs), intra-abdominal infections, wound infections, eye infections, musculoskeletal infections, bloodstream infections (bacteremia), skin infections, soft tissue infections, respiratory tract infections, post-surgical peritonitis, blood and brain infections	(32–35)
<i>E. sakazakii</i>	Neonatal meningitis, hydrocephalus, sepsis, and necrotizing enterocolitis.	(36–38)
<i>E. aerogenes</i>	Pneumonia, UTIs, bacteremia, wound infections, respiratory tract infections, skin infections, soft tissue infections, gastrointestinal infections, and septic infections, as well as fulminant forms of necrotizing meningitis in infants.	(39)

<i>Enterobacter</i>	Infection	Source
<i>E. hormaechei</i>	Bacteremia, lung infections, pneumonia, biliary tract infections, and UTIs	(40,41)
<i>E. amnigenus</i>	Endophthalmitis, bacteremia, UTIs	(42,43)
<i>E. asburiae</i>	Pneumonia, UTIs, intra-abdominal infections, and septicemia	
<i>E. carcinogenus</i>	Bacteremia and pneumonia	(44,45)
<i>E. cowanii</i>	Rhabdomyolysis, bacteremia, and acute cholecystitis	(46)
<i>E. dissolvans</i>	Septicemia and respiratory tract infections	(47)
<i>E. gergoviae</i>	Pneumonia, bacteremia, UTIs, respiratory tract infections, and peritonitis.	(48,49)
<i>E. kobei</i>	UTIs and sepsis	(50,51)
<i>E. ludwigii</i>	UTIs, bacteremia	(52,53)
<i>E. nimipressuralis</i>	Pseudobacteremia	(54)
<i>E. radicincitans</i>	Cholangiocarcinoma	(55)
<i>E. taylorae</i>	Bacteremia, UTIs, and meningitis.	(56)

The identified *Enterobacter* species exhibit extensive pathogenic potential, particularly as agents of opportunistic infections. *E. cloacae* is the predominant species, associated with a broad spectrum of infections including urinary tract infections, respiratory tract infections, bloodstream infections, and soft tissue infections. *E. sakazakii* (now classified as *Cronobacter sakazakii*) is notably linked to severe neonatal infections such as meningitis and sepsis. Other species, including *E. aerogenes*, *E. hormaechei*, and *E. gergoviae*, are frequently implicated in nosocomial infections such as pneumonia, bacteremia, and gastrointestinal infections. Less common species like *E. cowanii*, *E. dissolvans*, and *E. ludwigii* have been reported to cause serious conditions, including rhabdomyolysis, septicemia, and urinary tract infections. The detection of these species in ready-to-eat and street food environments underscores a tangible risk of infection transmission, especially in regions with inadequate sanitation.

DISCUSSION

Enterobacter Contamination Levels in Street Food in Various Countries

Foodborne infections are a serious health problem worldwide, affecting millions yearly (57). Globally, foodborne infections cause approximately 600 million cases and 420,000 deaths each year with the highest incidence in children under 5 years of age (3,58). This corresponds with the results in the United States, where foodborne pathogen contamination is expected to cause 9.4 million infections, 56,000 hospitalizations, and 1,350 deaths in 2022 (59). The United Kingdom reports approximately 2.4 million cases of foodborne infections each year with 180 deaths (60). The number of people suffering from foodborne diarrhea and other cases in developing countries exceeds 1.5 billion with 5 million deaths yearly (61). Therefore, the incidence suggests that the causative factor of infections refers to the microbiological quality of food. Various countries have provided alarming reports related to high pathogen contamination levels in food, specifically *Enterobacter*.

Sources of *Enterobacter* Contamination in Street Food

The human body is an ecological habitat for various microbial communities. *Enterobacter* is among the bacteria that can live on the skin and digestive tract of human, thereby playing a complex role in health and infections (62). Food poisoning, known as foodborne infections, is a condition caused by consuming food contaminated with microbes or toxins (63). The lack of effective implementation of regulations regarding safety and hygiene standards by sellers is a problem endangering consumer health (5). The main sources of bacterial contamination in food are generally caused by several factors, including unclean equipment, unhygienic sales locations, and inadequate sanitation facilities (64,65). Additionally, limited access to clean water, poor waste management, and storage of raw materials and food at inappropriate temperatures play a role in increasing contamination risk. These factors produce an environment supporting the growth and spread of bacteria, which can endanger consumer health. The method of preparing, processing, and serving food affects the microbiological quality (19).

Enterobacter contamination in street food primarily originates from raw materials used during food preparation, with *Enterobacter* species detected in water, wheat flour, meat, fruits, and vegetables (24,66–69). The equipment employed in street food processing is also a critical factor to consider. The hygiene practices of food handlers significantly influence the transmission of infections caused by *Enterobacter* bacteria. Moreover, *Enterobacter* can be transmitted directly or indirectly from the human body to food (70). Studies have demonstrated that food handlers may carry pathogenic *Enterobacteriaceae* on their hands and nails, which can lead to contamination during food preparation or handling (71–73).

Pathogenicity and *Enterobacter* Infection

Enterobacter is a genus of gram-negative, facultative anaerobic, rod-shaped, non-spore-forming, and motile bacteria belonging to *Enterobacteriaceae* family (74) known as opportunistic pathogens, which cause infections mainly in people with weakened immune systems and produce endotoxins and exotoxins contributing to virulence (33). This can be found in various habitats such as soil, plants, and the intestinal tract of humans and animals as well as in food (75). Furthermore, contamination of the bacteria in street food is transmitted through food handlers, cooking utensils, raw materials, water, and the environment (76,77). The presence of enterotoxigenic *Enterobacter* species in street food poses a significant public health risk, specifically due to the association with foodborne poisoning commonly referred to as foodborne infections. Currently, there are around 22 species belonging to the genus *Enterobacter*, namely *E. aerogenes*, *E. amnigenus*, *E. arachidis*, *E. asburiae*, *E. carcinogenus*, *E. cloacae*, *E. cowanii*, *E. dissolvans*, *E. gergoviae*, *E. helveticus*, *E. hormaechei*, *E. kobei*, *E. ludwigii*, *E. mori*, *E. nimipressuralis*, *E. oryzae*, *E. pulveris*, *E. pyrinus*, *E. radicincitans*, *E. soli*, *E. taylorae*, and *E. sakazakii* (33,78).

The pathogenicity and virulence factors of *Enterobacter spp.* are still not specifically known due to the lack of relevant studies. This group of bacteria has flagella playing a role in the pathogenicity which is used for motility/movement, biofilm formation, protein export, and adhesion (79). *Enterobacter* genus can excrete enterotoxins, alpha-hemolysin, and cytotoxins similar to Shiga-like toxins II and thiol-activated pore-forming cytotoxins (80,81). SLT-II inhibits protein synthesis and vascular damage, while thiol-activated pore-forming cytotoxins play a role in pore formation and cell lysis. In gram-negative bacteria, there is a type of secretion system (T3SS) that functions to inject effector proteins into eukaryotic host cells, where this system plays a role in the virulence of pathogenic bacteria (82). *E. cloacae* has 2 types of type VI secretion systems (T6SS), namely T6SS-1 and T6SS-2, which enhances the ability of bacteria to survive in a competitive environment (83). *E. cloacae* is capable of attaching to and penetrating epithelial cells as well as inducing apoptosis in Hep2 cells, which could be a crucial mechanism contributing to the development of infections (84). *E. cloacae complex* has Pqc plasmids containing virulence-encoding genes (ter and sea) and resistance-encoding genes (blaCTX-M-9, qnrA1, aadB, aadA2, sukK, and sat), contributing to the virulence and adaptation of *E. cloacae*¹. This complex comprises *E. cloacae*, *E. asburiae*, *E. hormaechei*, *E. kobei*, *E. ludwigii*, *E. mori*, and *E. nimipressuralis*, grouped based on the similarity of approximately 60% of the genomes (85). Additionally, *E. hormaechei* bacteria are more dangerous due to being an EHOS (*E. hormaechei* Outbreak Strain) containing a High Pathogenicity Island (HPI) (86). A key genetic element in pathogenicity known as HPI encodes virulence factors essential for the acquisition, regulation, and transport of iron, promoting pathogenicity and the ability of bacteria to cause infections (87,88). The ability to assimilate iron through chelators is crucial for bacterial metabolism and the establishment of infection. Siderophore encoding genes are commonly observed in HPI, particularly in *Yersinia spp.* Among these genes, the *irp2* gene has been identified in *Enterobacter spp.* (89). Moreover, *E. cloacae* can express Curli fimbriae, which are extracellular protein fibers included in the amyloid category, functioning to increase bacterial adhesion to surfaces, cell aggregation, and biofilm formation (90). The genes encoding Curli are grouped into two operons, namely *csgBA(C)* and *csgDEFG*, with studies showing that 11 (78.6%) of 14 isolates has the *csgBA* operon. The ability of *E. cloacae* isolates to form biofilms was significantly correlated with the mRNA expression levels of the *csgA* encoding Curli subunit and *csgD* activating the *csgBA* operon promoter as well as *csgE*, *csgF*, and *csgG*. Differences in pathogenicity between *E. aerogenes* and *E. cloacae* have been reported in various investigations (91). *E. aerogenes* has virulence-encoding genes *fimH* and *mrkD* such as those encoding fimbrial adhesins types 1 and 3, with a key role in adhesion and biofilm

formation, which are crucial aspects of bacterial virulence (33,91,92). Furthermore, iron transport genes *kfu*, *entB*, and *ybtS* participating in the production of siderophores were identified in *E. aerogenes*, which facilitate iron absorption through the production of siderophores such as enterobactin and yersiniabactin. These genes enhance the ability of bacteria to survive and reproduce in competitive host environments (91). *Enterobacter aerogenes* can produce beta-lactamases, specifically Extended-Spectrum Beta-Lactamases (ESBLs) and AmpC cephalosporinase playing a role in virulence and adaptability of resistance genes (93,94).

Strategy and Recommendations for Addressing *Enterobacter* Contamination of Street Food in Developing Countries

Contamination of *Enterobacter* spp. in street food poses a serious public health challenge in developing countries, contributing substantially to the burden of microbiological infectious diseases, particularly among vulnerable populations. Therefore, the development of effective management strategies should adopt a multidisciplinary approach integrating education, regulation, technology, and enhanced infrastructure.

First, empowering street food vendors through education and training programs focused on food hygiene and sanitation principles is essential. Studies indicate that vendors' lack of knowledge regarding sanitation practices, safe food processing, and raw material handling are major contributors to the risk of *Enterobacter* spp. contamination (95). Continuous training should be implemented promptly to enhance vendors' competencies in controlling microbial contamination, emphasizing effective handwashing, guaranteed use of clean water, and standardized food processing and storage methods.

Furthermore, the implementation of strict and well-documented hygiene and sanitation standard operating procedures (SOPs) is necessary to reduce variability in hygiene practices in the field (96). SOPs should cover cleaning of cooking utensils, work surfaces, waste management, and handling of raw and cooked foods, tailored specifically to the characteristics of local street food. The enforcement of these SOPs requires regular internal and external audits to ensure compliance and effectiveness.

Government regulation plays a crucial role in supporting these interventions. Local authorities need to strengthen supervision, certification, and law enforcement related to street food safety (97). Systematic, risk-based periodic inspections should be conducted to identify and mitigate critical contamination points. Additionally, awarding hygiene certificates to vendors who meet safety standards can serve as an incentive, motivating businesses to adopt safe and hygienic practices. The availability and quality of clean water is another critical factor demanding serious attention. Since water is extensively used for washing materials and equipment, improving access to microbiologically safe water sources and implementing water treatment systems at street food vending sites are essential to eliminate potential contaminants.

Moreover, consumer empowerment through education about the health risks associated with consuming unhygienic food is vital. Increased consumer awareness of food safety encourages the selection of safer food products, thereby motivating vendors to improve their food handling practices. Continuous monitoring to track the presence of *Enterobacter* spp. in street food, contamination pathways, and antibiotic resistance patterns is imperative. This monitoring supports the formulation of adaptive and effective control policies and provides a basis for evaluating the success of intervention programs.

In summary, addressing *Enterobacter* spp. contamination in street food in developing countries requires multisectoral collaboration and an evidence-based approach to achieve optimal food safety and reduce the risk of microbiological infectious diseases in the community.

CONCLUSION

Street food in various developing countries has been found to be contaminated with *Enterobacter* at levels that pose significant public health risks. The prevalence of *Enterobacter* contamination in street food varies widely, ranging from 5.36% up to 100%. *Enterobacter aerogenes* was identified as the most dominant species, with the highest prevalence detected in ready-to-eat (RTE) cooked rice samples from Nigeria. Sources of *Enterobacter* contamination in street food include raw materials, cooking utensils,

water, and food handlers, with prevalence rates ranging from 0.99% to 100%. Wheat flour was recorded as the raw material with the highest contamination level, where *E. sakazakii* prevalence reached 100%. The high pathogenicity, virulence, and antibiotic resistance capabilities of *Enterobacter* make it a significant causative agent in various infections. The predominant bacterial species, *E. sakazakii* and *E. cloacae*, are frequently associated with urinary tract infections, meningitis, gastrointestinal infections, pneumonia, septicemia, and bacteremia.

This literature review underscores the critical need for multidisciplinary prevention and control strategies to mitigate *Enterobacter* contamination in street food and reduce the associated public health risks. Recommended strategies include the implementation of good hygiene and sanitation practices, the use of safe raw materials, education and training for food vendors, and regular food safety monitoring. Furthermore, additional research is warranted to elucidate contamination mechanisms, characterize virulence factors, and understand antibiotic resistance patterns to support the development of more effective policies and interventions.

FUNDING

This research received no external funding

CONFLICTS OF INTEREST

The authors declare no conflict of interest

REFERENCES

1. Mengistu DA, Belami DD, Tefera AA, Alemeshet Asefa Y. Bacteriological quality and public health risk of ready-to-eat foods in developing countries: Systematic review and meta analysis. *Microbiology Insights*. 2022 Jan;15:117863612211139.
2. Havelaar AH, Kirk MD, Torgerson PR, Gibb HJ, Hald T, Lake RJ, et al. World Health Organization global estimates and regional comparisons of the burden of foodborne disease in 2010. Von Seidlein L, editor. *PLoS Med*. 2015 Dec 3;12(12):e1001923.
3. Lee H, Yoon Y. Etiological agents implicated in foodborne illness world wide. *Food Sci Anim Resour*. 2021 jan;41(1):1–7.
4. Barreira MJ, Marcos S, Flores CV, Lopes TT, Moura IB, Correia CB, et al. Microbiological quality of ready-to-eat street foods in Lisbon, Portugal. *Discov Food*. 2024 Jun 24;4(1):45.
5. Sabuj AAM, Haque ZF, Younus MdI, Pondit A, Barua N, Hossain MdG, et al. Microbial risk assessment of ready-to-eat fast foods from different street-vended restaurants. *Int J One Health*. 2020;6(1):41–8.
6. Lyytinen OL, Dapuliga C, Wallinger D, Patpatia S, Audu BJ, Kiljunen SJ. Three novel *Enterobacter cloacae* bacteriophages for therapeutic use from Ghanaian natural waters. *Arch Virol*. 2024 Aug;169(8):156.
7. Herawati H, Sakati SN, Sumarto Z. Kualitas bakteriologis pada peralatan Makan di Warung Makan Kadompe di Kota Luwuk Kabupaten Banggai. *JIK*. 2022 Aug 29;16(2):200–6.
8. Asiegbu CV, Lebelo SL, Tabit FT. Microbial quality of ready-to-eat street vended food groups sold in the Johannesburg Metropolis, South Africa. *JFQHC [Internet]*. 2020 Feb 27 [cited 2024 Oct 7]; Available from: <https://publish.kne-publishing.com/index.php/JFQHC/article/view/2448>
9. Hossain MB, Mahbub NDB, Chowdhury MMK, Rahaman MM. Molecular characterization of *Enterobacter* and *Escherichia coli* pathotypes prevalent in the popular street foods of Dhaka City and their multidrug resistance. *Bangla J Microbiol*. 2019 Jan 1;34(2):67–72.
10. Dela H, Egyir B, Behene E, Sulemana H, Tagoe R, Bentil R, et al. Microbiological quality and antimicrobial resistance of bacteria species recovered from ready-to-eat food, water samples, and palm swabs of food vendors in Accra, Ghana. *International Journal of Food Microbiology*. 2023 Jul;396:110195.
11. Izevbuwa O, Okhuebor S. Microbiological assessment of ready to eat food from selected street vending food locations in Ikpoba-okha local government area of Edo State. *BE*. 2021 Jan 20;4(1):20–4.

12. Sonune NA. Assessment of bacteriological quality of street food and their antibiotic profiling. SAR J Pathol Microbiol. 2022 Jul 13;3(4):39–45.
13. Amare A, Worku T, Ashagirie B, Adugna M, Getaneh A, Dagne M. Bacteriological profile, antimicrobial susceptibility patterns of the isolates among street vended foods and hygienic practice of vendors in Gondar town, Northwest Ethiopia: a cross sectional study. BMC Microbiol. 2019 Dec;19(1):120.
14. Murase K. Cytolysin A (ClyA): A Bacterial virulence factor with potential applications in nanopore technology, vaccine development, and tumor therapy. Toxins. 2022 Jan 21;14(2):78.
15. Hassanin F, Hassan M, Helmy S. *Enterobacteriaceae* in meat products sold by street vendors. Benha Veterinary Medical Journal. 2015 Dec 1;29(2):74–9.
16. Igbinosa IH, Beshiru A, Egharevba NE, Igbinosa EO. Distribution of *Enterobacteria* in Ready-to-eat food in cafeterias and retail food outlets in Benin City: Public health implications. j com med and phc. 2020 aug 28;32(2):80–94.
17. Wang S, Xiao SZ, Gu FF, Tang J, Guo XK, Ni YX, et al. Antimicrobial susceptibility and molecular epidemiology of clinical *Enterobacter cloacae* bloodstream isolates in Shanghai, China. Galdiero M, editor. PLoS ONE. 2017 Dec 15;12(12):e0189713.
18. Ogunyemi A, Buraimoh O, Onuorah N, Ezeugwu S, Odetunde S, Olumuyiwa E. Bacteria associated with contamination of ready-to-eat (RTE) cooked rice in Lagos-Nigeria. Int J Bio Chem Sci. 2016 Feb 5;9(5):2324.
19. Nyenje ME, Odjadjare CE, Tanih NF, Green E, Ndip RN. Foodborne pathogens recovered from ready-to-eat foods from roadside cafeterias and retail outlets in Alice, Eastern Cape Province, South Africa: Public health implications. IJERPH. 2012 Jul 27;9(8):2608–19.
20. Sharma I, Mazumdar J. Assessment of bacteriological quality of ready to eat food vended in streets of Silchar city, Assam, India. Indian Journal of Medical Microbiology. 2014 Apr;32(2):169–71.
21. Budiarto TY, Amarantini C, Prihatmo G, Restiani R, Putri Y, Kindagen V, et al. Detection of coliforms and enteric pathogens in favorite snack food sold in Yogyakarta City: In Yogyakarta, Indonesia; 2021 [cited 2024 Oct 15]. Available from: <https://www.atlantispress.com/article/125953691>
22. Lou X, Si G, Yu H, Qi J, Liu T, Fang Z. Possible reservoir and routes of transmission of *Cronobacter* (*Enterobacter sakazakii*) via wheat flour. Food Control. 2014 Sep;43:258–62.
23. Razzaq R, Farzana K, Mahmood S, Murtaza G. Microbiological analysis of street vended vegetables in Multan City, Pakistan: A Public Health Concern.
24. Falomir MP, Rico H, Gozalbo D. *Enterobacter* and *Klebsiella* species isolated from fresh vegetables marketed in Valencia (Spain) and their clinically relevant resistances to chemotherapeutic agents. Foodborne Pathogens and Disease. 2013 Dec;10(12):1002–7.
25. Oranusi S, Olorunfemi OJ. Microbiological safety evaluation of street vended ready-to-eat fruits sold in Ota, Ogun state, Nigeria.
26. Attien P, Dadie T, Sina H, Kouassi CK, Baba-Moussa L. Sanitary risk factors and microbial profile identification by MALDI-TOF of street ready-to-eat meat products contaminants. Int J Res Granthaalayah. 2020 Jun 30;5(12):164–75.
27. Shiningeni D, Chimwamurombe P, Shilangale R, Misihairabgwi J. Prevalence of pathogenic bacteria in street vended ready-to-eat meats in Windhoek, Namibia. Meat Science. 2019 Feb;148:223–8.
28. Saksena R, Malik M, Gaiind R. Bacterial contamination and prevalence of antimicrobial resistance phenotypes in raw fruits and vegetables sold in Delhi, India. Journal of Food Safety. 2020 Feb;40(1):e12739.
29. Shayeghi F, Matini E, Rahbar N, Mojri N, Hosseini SS, Abdollahi M, et al. Microbial contamination of kitchen instruments as a minatory to human health.
30. Haymanot Tasew TA. Contamination of bacteria and associated factors among food handlers working in the student cafeterias of Jimma University main campus, Jimma, South West Ethiopia. Altern Integ Med [Internet]. 2015 [cited 2024 Oct 17];04(01). Available from: <http://www.esciencecentral.org/journals/contamination-of-bacteria-and-associated-factors-among-food-handlers-working-in-the-student-cafeterias-of-jimma-university-main-campus-jimma-south-west-ethiopia-2327-5162.1000185.php?aid=40793>

31. Aruwa CE, Ogundare O. Microbiological Quality assessment of pupuru and plantain flours in an urban market in Akure, Ondo State, South Western Nigeria. *OALib*. 2017;04(08):1–11.
32. Ceronifrom D. Musculoskeletal postoperative infections due to *Enterobacter cloacae* complex: A new reality? *J Med Microb Diagn* [Internet]. 2014 [cited 2024 Oct 29];03(01). Available from: <https://www.omicsonline.org/open-access/musculoskeletal-postoperative-infections-due-to-Enterobacter-cloacaecomplex-a-new-reality-2161-0703.1000128.php?aid=21218>
33. Davin-Regli A, Lavigne JP, Pagès JM. *Enterobacter* spp.: Update on taxonomy, clinical aspects, and emerging antimicrobial resistance. *Clin Microbiol Rev*. 2019 Sep 18;32(4):e00002–19.
34. Mbula J, Abdulsalami MS, Vantsawa PA, Onuh K, Ozojiofor UO, Ayodele SB, et al. Isolation and molecular identification of *Enterobacter cloacae* from pregnant women with urinary tract infections in Biu, North Eastern Nigeria, Borno State. *JAMB*. 2023 Apr 18;23(5):1–15.
35. Mezzatesta ML, Gona F, Stefani S. *Enterobacter cloacae* complex: Clinical impact and emerging antibiotic resistance. *Future Microbiol*. 2012 Jul;7(7):887–902.
36. Šín R, Štruncová D, Čechurová L. Clinical picture, diagnostics and treatment of bacterial meningitis.
37. Yamamoto H, Miyamoto Y, Yamamoto H. A case of bacterial meningitis with burst waves of local onset on ictal Electroencephalography. *Pediatrics International*. 2019 Dec;61(12):1263–4.
38. Zhou X, Fu S, Gao J, Chen H. *Enterobacter sakazakii*: an emerging foodborne pathogenic bacterium. *Ann Microbiol*. 2012 Mar;62(1):1–5.
39. Rizi KS. Clinical and pathogenesis overview of *Enterobacter* infections. 6(4).
40. Appleberry H, Patel R, Singh K, Wolfe AJ, Putonti C, Kula A. Draft genomes of two *Enterobacter hormaechei* strains isolated from catheterized urine samples from females experiencing overactive bladder symptoms. Klepac-Ceraj V, editor. *Microbiol Resour Announc*. 2024 Aug 13;13(8):e00491–24.
41. Jiménez-Castillo RA, Aguilar-Rivera LR, Carrizales-Sepúlveda EF, Gómez-Quiroz RA, Llantada-López AR, González-Aguirre JE, et al. A case of round pneumonia due to *Enterobacter hormaechei*: the need for a standardized diagnosis and treatment approach in adults. *Rev Inst Med trop S Paulo*. 2021;63:e3.
42. Mohammed MA, Alnour TMS, Shakurfo OM, Aburass MM. Prevalence and antimicrobial resistance pattern of bacterial strains isolated from patients with urinary tract infection in Messalata Central Hospital, Libya. *Asian Pacific Journal of Tropical Medicine*. 2016 Aug;9(8):771–6.
43. Westerfeld C, Papaliodis GN, Behlau I, Durand ML, Sobrin L. *Enterobacter amnigenus* endophthalmitis. *Retinal Cases & Brief Reports*. 2009;3(4):409–11.
44. Cruz R, López E, Meneses C, Alegría Y. Bacteriemia por *Enterobacter cancerogenus* en un paciente con trauma pélvico. *Rev chil infectol*. 2022 Apr;39(2):218–20.
45. Demir T, Baran G, Buyukguclu T, Sezgin FM, Kaymaz H. Pneumonia due to *Enterobacter cancerogenus* infection. *Folia Microbiol*. 2014 Nov;59(6):527–30.
46. Berinson B, Bellon E, Christner M, Both A, Aepfelbacher M, Rohde H. Identification of *Kosakonia cowanii* as a rare cause of acute cholecystitis: Case report and review of the literature. *BMC Infect Dis*. 2020 Dec;20(1):366.
47. Chen Q, Lin Y, Li Z, Lu L, Li P, Wang K, et al. Characterization of a new transposon, Tn6696, on a blaNDM-1-Carrying plasmid from multidrug-resistant *Enterobacter cloacae* ssp. *dissolvens* in China. *Front Microbiol*. 2020 Sep 15;11:525479.
48. Kesieme EB, Kesieme CN, Akpede GO, Okonta KE, Dongo AE, Gbolagade AM, et al. Tension pneumatocele due to *Enterobacter gergoviae* pneumonia: A case report. *Case Reports in Medicine*. 2012;2012:1–3.
49. Satlin MJ, Jenkins SG, Chen L, Helfgott D, Feldman EJ, Kreiswirth BN, et al. Septic shock caused by *Klebsiella pneumoniae* carbapenemase-producing *Enterobacter gergoviae* in a neutropenic patient with leukemia. *J Clin Microbiol*. 2013 Aug;51(8):2794–6.
50. Manandhar S, Nguyen Q, Pham DT, Amatya P, Rabaa M, Dongol S, et al. A fatal outbreak of neonatal sepsis caused by mcr-10-carrying *Enterobacter kobei* in a tertiary care hospital in Nepal. *Journal of Hospital Infection*. 2022 Jul;125:60–6.

51. Zeng H, Tan Y, Su J, Gao F, Lei T, Liang B. Co-occurrence of blaNDM-1, rmtC, and mcr-9 in multidrug-resistant *Enterobacter kobei* strain isolated from an infant with urinary tract infection. *Journal of Global Antimicrobial Resistance*. 2023 Jun;33:221–6.
52. Shafeeq S, Wang X, Lünsdorf H, Brauner A, Römling U. Draft genome sequence of the urinary catheter isolate *Enterobacter ludwigii* CEB04 with high biofilm forming capacity. *Microorganisms*. 2020 Apr 5;8(4):522.
53. Wagner L, Bloos F, Vylkova S. Bloodstream infection due to *Enterobacter ludwigii*, correlating with massive aggregation on the surface of a central venous catheter. *Infection*. 2020 Dec;48(6):955–8.
54. Kim DM, Jang SJ, Neupane GP, Jang MS, Kwon SH, Kim SW, et al. *Enterobacter nimipressuralis* as a cause of pseudobacteremia. *BMC Infect Dis*. 2010 Dec;10(1):315.
55. Mertschnigg T, Patz S, Becker M, Feierl G, Ruppel S, Bunk B, et al. First report of *Kosakonia radicincitans* bacteraemia from Europe (Austria) - Identification and whole-genome sequencing of strain DSM 107547. *Sci Rep*. 2020 Feb 6;10(1):1948.
56. Rubinstien EM, Klevjer-Anderson P, Smith CA, Drouin MT, Patterson JE. *Enterobacter taylorae*, a new opportunistic pathogen: Report of four cases. *J Clin Microbiol*. 1993 Feb;31(2):249–54.
57. De Vogli R, Kouvonen A, Gimeno D. The influence of market deregulation on fast food consumption and body mass index: A cross-national time series analysis. *Bull World Health Organ*. 2014 Feb 1;92(2):99-107A.
58. Verma R, Mishra S. Food safety issues of street foods and dietary practices by school going adolescents.
59. Weller DL, Ray LC, Payne DC, Griffin PM, Hoekstra RM, Rose EB, et al. An enhanced method for calculating trends in infections caused by pathogens transmitted commonly through food [Internet]. *Epidemiology*; 2022 [cited 2024 Oct 11]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2022.09.14.22279742>
60. Holland D, Thomson L, Mahmoudzadeh N, Khaled A. Estimating deaths from foodborne disease in the UK for 11 key pathogens. *BMJ Open Gastroenterol*. 2020 Jun;7(1):e000377.
61. Majeed A. Food toxicity: Contamination sources, health implications and prevention.
62. Micheel V, Hogan B, Rakotoarivelo R, Rakotozandrindrainy R, Razafimanatsoa F, Razafindrabe T, et al. Identification of nasal colonization with β -lactamase-producing *Enterobacteriaceae* in patients, health care workers and students in Madagascar. *European Journal of Microbiology and Immunology*. 2015 Mar;5(1):116–25.
63. Mehboob A, Abbas T. Evaluation of microbial quality of street food in Karachi City, Pakistan: An epidemiological study. *Microbiol Res (Pavia)* [Internet]. 2019 Jan 29 [cited 2024 Oct 2];10(1). Available from: <https://www.pagepress.org/journals/index.php/mr/article/view/7463>
64. Asiegbu CV, Lebelo SL, Tabit FT. The food safety knowledge and microbial hazards awareness of consumers of ready-to-eat street-vended food. *Food Control*. 2016 Feb;60:422–9.
65. Raza J, Asmat TM, Mustafa MZ, Ishtiaq H, Mumtaz K, Jalees MM, et al. Contamination of ready-to-eat street food in Pakistan with *Salmonella* spp.: Implications for consumers and food safety. *International Journal of Infectious Diseases*. 2021 May;106:123–7.
66. Akinde SB, Sunday AA, Adeyemi FM, Fakayode IB, Oluwajide OO, Adebunmi AA, et al. Microbes in irrigation water and fresh vegetables: Potential pathogenic bacteria assessment and implications for food safety. *Appl Biosaf*. 2016 Jun;21(2):89–97.
67. Mladenović KG, Grujović MŽ, Kiš M, Furmeg S, Tkalec VJ, Stefanović OD, et al. *Enterobacteriaceae* in food safety with an emphasis on raw milk and meat. *Appl Microbiol Biotechnol*. 2021 Dec;105(23):8615–27.
68. Pintor-Cora A, Alegría Á, Ramos-Vivas J, García-López ML, Santos JA, Rodríguez-Calleja JM. Antimicrobial-resistant *Enterobacter cloacae* complex strains isolated from fresh vegetables intended for raw consumption and their farm environments in the Northwest of Spain. *LWT*. 2023 Oct;188:115382.
69. Umar Mustapha M, Halimoon N, Wan Johari WL, Abd Shukor MohdY. Enhanced carbofuran degradation using immobilized and free cells of *Enterobacter* sp. Isolated from Soil. *Molecules*. 2020 Jun 16;25(12):2771.

70. Amare A, Eshetie S, Kasew D, Moges F. High prevalence of fecal carriage of extended-spectrum beta-lactamase and carbapenemase-producing *Enterobacteriaceae* among food handlers at the University of Gondar, Northwest Ethiopia. Algammal AM, editor. PLoS ONE. 2022 Mar 17;17(3):e0264818.
71. Khurana KM, Hassani US. Presence of pathogenic *Enterobacteriaceae* on fingertips of food handlers. Journal of Datta Meghe Institute of Medical Sciences University. 2024 Apr;19(2):281–5.
72. Moges Tiruneh MD. Bacterial Profile and antimicrobial susceptibility pattern among food handlers at Gondar University Cafeteria, Northwest Ethiopia. J Infect Dis Ther [Internet]. 2013 [cited 2024 Oct 17];01(02). Available from: <http://www.esciencecentral.org/journals/bacterial-profile-and-antimicrobial-susceptibility-pattern-among-food-handlers-at-gondar-university-cafeteria-northwest-ethiopia-2332-0877.1000105.php?aid=14902>
73. Zagloul D, Khodari Y, Othman RAM, Farooq M. Prevalence of intestinal parasites and bacteria among food handlers in a tertiary care hospital. Niger Med J. 2011;52(4):266.
74. Bolourchi N, Fereshteh S, Noori Goodarzi N, Badmasti F. Subtractive genomic analysis for computational identification of putative immunogenic targets against clinical *Enterobacter cloacae* complex. Marsano RM, editor. PLoS ONE. 2022 Oct 13;17(10):e0275749.
75. Ben Said L, Jouini A, Klibi N, Dziri R, Alonso CA, Boudabous A, et al. Detection of extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae* in vegetables, soil and water of the farm environment in Tunisia. International Journal of Food Microbiology. 2015 Jun;203:86–92.
76. Nisa TT, Sugawara Y, Hamaguchi S, Takeuchi D, Abe R, Kuroda E, et al. Genomic characterization of carbapenemase-producing *Enterobacterales* from Dhaka food markets unveils the spread of high-risk antimicrobial-resistant clones and plasmids co-carrying *bla* NDM and *mcr-1.1*. JAC-Antimicrobial Resistance. 2024 Jul 3;6(4):dlae124.
77. Rane S. Street vended food in developing world: Hazard analyses. Indian J Microbiol. 2011 Jan;51(1):100–6.
78. Grimont F, Grimont PAD. The genus *Enterobacter*. In: Dworkin M, Falkow S, Rosenberg E, Schleifer KH, Stackebrandt E, editors. The Prokaryotes [Internet]. New York, NY: Springer New York; 2006 [cited 2024 Oct 31]. p. 197–214. Available from: http://link.springer.com/10.1007/0-387-30746-X_9
79. Haiko J, Westerlund-Wikström B. The role of the bacterial flagellum in adhesion and virulence. Biology. 2013 Oct 25;2(4):1242–67.
80. Barnes AI, Ortiz C, Paraje MG, Balanzino LE, Albasa I. Purification and characterization of a cytotoxin from *Enterobacter cloacae*. Can J Microbiol. 1997 Aug 1;43(8):729–33.
81. Krzysińska S, Mokracka J, Koczura R, Kaznowski A. Cytotoxic activity of *Enterobacter cloacae* human isolates. FEMS Immunol Med Microbiol. 2009 Aug;56(3):248–52.
82. Horna G, Ruiz J. Type 3 secretion system of *Pseudomonas aeruginosa*. Microbiological Research. 2021 May;246:126719.
83. Hernández-Martínez G, Ares MA, Rosales-Reyes R, Soria-Bustos J, Yañez-Santos JA, Cedillo ML, et al. The nucleoid protein HU positively regulates the expression of type VI secretion systems in *Enterobacter cloacae*. D'Orazio SEF, editor. mSphere. 2024 May 29;9(5):e00060-24.
84. Krzysińska S, Koczura R, Mokracka J, Puton T, Kaznowski A. Isolates of the *Enterobacter cloacae* complex induce apoptosis of human intestinal epithelial cells. Microbial Pathogenesis. 2010 Sep;49(3):83–9.
85. Paauw A, Caspers MPM, Schuren FHJ, Leverstein-van Hall MA, Delétoile A, Montijn RC, et al. Genomic diversity within the *Enterobacter cloacae* Complex. Redfield RJ, editor. PLoS ONE. 2008 Aug 21;3(8):e3018.
86. Paauw A, Caspers MPM, Leverstein-van Hall MA, Schuren FHJ, Montijn RC, Verhoef J, et al. Identification of resistance and virulence factors in an epidemic *Enterobacter hormaechei* outbreak strain. Microbiology. 2009 May 1;155(5):1478–88.
87. Royer G, Clermont O, Marin J, Condamine B, Dion S, Blanquart F, et al. Epistatic interactions between the high pathogenicity island and other iron uptake systems shape *Escherichia coli* extra-intestinal virulence. Nat Commun. 2023 Jun 20;14(1):3667.

88. Shan C, Liu C, Lu Q, Fu G, Shah SAH, Akhtar RW, et al. High pathogenicity island (HPI) main structural genes and their bioinformatics analysis in clinically isolated *E. coli* from Saba Pigs. PJZ [Internet]. 2022 [cited 2024 Nov 1];55(2). Available from: <http://researcherslinks.com/current-issues/High-Pathogenicity-Island-HPI-Main-Structural-Genes/20/1/5674/html>
89. Souza Lopes AC, Rodrigues JF, Cabral AB, Da Silva ME, Leal NC, Da Silveira VM, et al. Occurrence and analysis of *irp2* virulence gene in isolates of *Klebsiella pneumoniae* and *Enterobacter* spp. from microbiota and hospital and community-acquired infections. Microbial Pathogenesis. 2016 Jul;96:15–9.
90. Kim SM, Lee HW, Choi YW, Kim SH, Lee JC, Lee YC, et al. Involvement of curli fimbriae in the biofilm formation of *Enterobacter cloacae*. J Microbiol. 2012 Feb;50(1):175–8.
91. Azevedo PAA, Furlan JPR, Oliveira-Silva M, Nakamura-Silva R, Gomes CN, Costa KRC, et al. Detection of virulence and β -lactamase encoding genes in *Enterobacter aerogenes* and *Enterobacter cloacae* clinical isolates from Brazil. Brazilian Journal of Microbiology. 2018 Nov;49:224–8.
92. El Fertat-Aissani R, Messai Y, Alouache S, Bakour R. Virulence profiles and antibiotic susceptibility patterns of *Klebsiella pneumoniae* strains isolated from different clinical specimens. Pathologie Biologie. 2013 Oct;61(5):209–16.
93. Claeys G, De Baere T, Wauters G, Vandecandelaere P, Verschraegen G, Muylaert A, et al. Extended-spectrum β -lactamase (ESBL) producing *Enterobacter aerogenes* phenotypically misidentified as *Klebsiella pneumoniae* or *K. terrigena*. BMC Microbiol. 2004 Dec 24;4(1):49.
94. Davin-Regli A, PagÃ’s JM. *Enterobacter aerogenes* and *Enterobacter cloacae*; versatile bacterial pathogens confronting antibiotic treatment. Front Microbiol [Internet]. 2015 May 18 [cited 2024 Nov 1];6. Available from: http://www.frontiersin.org/Antimicrobials%2c_Resistance_and_Chemotherapy/10.3389/fmicb.2015.00392/abstract
95. Madilo FK, Islam MN, Letsyo E, Roy N, Klutse CM, Quansah E, et al. Foodborne pathogens awareness and food safety knowledge of street-vended food consumers: A case of university students in Ghana. Heliyon. 2023 Jul;9(7):e17795.
96. Jambre KGE, Lagorra MJP. Compliance to sanitation standard operating procedures: An evaluation of street food handlers in Dipolog City, Philippines. 2022;(6).
97. Huynh-Van B, Vuong-Thao V, Huynh-Thi-Thanh T, Dang-Xuan S, Huynh-Van T, Tran-To L, et al. Factors associated with food safety compliance among street food vendors in Can Tho city, Vietnam: implications for intervention activity design and implementation. BMC Public Health. 2022 Dec;22(1):94.